

THERAPEUTIC EFFICACY OF ATROPINE AGAINST CHOLINESTERASE INHIBITORS TOXICITY IN CHICKS STRESSED WITH HYDROGEN PEROXIDE

Yaareb J. Mousa^{*}, Muna H. I. Al-Zubaidy^{*}, Sawsen M. Ameen^{*} and Fouad K.
Mohammad^{**}

^{*} Department of Physiology, Biochemistry and Pharmacology, College of Veterinary
Medicine, University of Mosul, Mosul, Iraq.

^{**} Ministry of Higher Education and Scientific Research, Baghdad, Iraq.

(Received 15 May 2016 ,Accepted 26 may 2016)

Keywords: Atropine, Oxidative Stress, H₂O₂, Chicks.

ABSTRACT

The aim of this study was to evaluate therapeutic efficacy of atropine in case of cholinesterase (ChE) inhibitors toxicity in chicks subjected to oxidative stress (OS) with hydrogen peroxide (H₂O₂). H₂O₂ at 0.5 % in drinking water induced OS at age of 7-14 days in the chicks when given each day for 14 consecutive days. There was no change in the acute median lethal dose (LD₅₀) for diazinon (7.9 mg/kg, orally) for both H₂O (control) and H₂O₂ groups. The therapeutic efficacy of atropine was decreased (265 %) when measuring the median effective dose (ED₅₀) for treating the diazinon toxicity in H₂O group (2.6 mg/kg, i.m.) and in the H₂O₂ group (9.5 mg/kg, i.m.). The signs of acute diazinon toxicity were found to increase in H₂O₂ groups when compared to H₂O group and there is slight decrease in plasma and whole brain ChE activities in H₂O₂ group when when compared to H₂O group. The data revealed a decline in atropine efficacy for

treatment diazinon toxicity in chicks that suffered from OS status and it is recommended here to increase the dose of atropine in this case.

INTRODUCTION

Atropine is the drug of choice which is used as an antidote for the treatment of cholinesterase (ChE) inhibitors toxicity such as organophosphates and carbamates in human and animals by blocking the muscarinic receptors, thereby diminishing the excess acetylcholine from binding to it (1-3). Diazinon is an organophosphate insecticides that induces toxicity through inhibiting acetylcholinesterase in living tissues (2,3). Studies have recommended using the powerful oxidant agent hydrogen peroxide (H₂O₂) for induction of OS in laboratory animals such rats and chicks (4-7). It was found that OS modulates the pharmacological response to drugs in chicks and if H₂O₂ was given at 0.5 % in drinking water, it induces OS in 7-14 day-old chicks when given fresh each day (5-7). H₂O₂ induces OS through its capacity to increase the reactive oxygen species (ROS) and free radicals formation in the tissues (8,9). The aim of the present study was to evaluate the therapeutic efficacy of atropine in case of ChE inhibitors (diazinon as a model) toxicity in chicks suffered from OS induced by H₂O₂ since no previous studies dealt with such a life threatening situation.

MATERIALS AND METHODS

Experimental animals and induction of OS

Day-old broiler chicks of both sexes (body weight 45-125 g) were used. They were kept in animal house at a temperature of 32–35 °C with constant lighting, and wood shavings as floor litter. The chicks had free access to drinking water and feed along the

experiment. The Scientific Committee of the College of Veterinary Medicine at the University of Mosul reviewed and approved the protocol of this study. To induce OS, day-old chicks were either provided with tap water (H₂O-control group) or with fresh H₂O₂ (Thomas Baker Chemical Ltd., U.K.) in tap water as 0.5% v/v each day for two weeks in order to produce an OS as reported in previous studies (4,5,7).

Determination of acute median lethal dose (LD₅₀) of diazinon in H₂O (control) and H₂O₂ (stressed) groups

The acute LD₅₀ of diazinon (60%, VAPCO, Jordan) was measured for both the control and stressed groups according to the up-and-down method described before (10). The initial dose of diazinon was 7 mg/kg, orally for both groups. The onset and signs of acute toxicity of diazinon in the chicks during 24 hours were also recorded in this experiment.

Measuring the median effective dose (ED₅₀) of atropine in the control and stressed groups for alteration of acute diazinon toxicity

After determination of the acute LD₅₀ value of diazinon (7.9 mg/kg, orally), the dose of diazinon (10 mg/kg, orally) was chosen that causes acute toxicity leading to lethality in the chicks. The ED₅₀ of atropine (1%, Alsharq for Veterinary Drugs, Syria) was determined for both the control and stressed groups according to up-and-down method (10). The onset and signs of acute toxicity of diazinon during 24 hours were measured to evaluate the differences between these two groups.

Therapeutic efficacy of atropine against acute diazinon toxicity in control and stressed chicks with H₂O₂

Atropine was injected at 6 mg/kg, i.m. (this dose was chosen according to ED₅₀ value of atropine) immediately after diazinon dosing at 10 mg/kg, orally for both the control and stressed chicks (6 chicks/group). The onset of acute diazinon toxicity, toxic signs and toxicity scores were evaluated for both groups.

In vivo determination of ChE activity in plasma and whole brain of the control and stressed chicks

Atropine was injected at 6 mg/kg, i.m. immediately after diazinon dosing at 10 mg/kg, orally for both the control and stressed chicks (6 chicks/group). The chicks were euthanized and blood samples and whole brain were collected after 2 hours of atropine and diazinon administration. The ChE activity was determined according to a modified electrometric method described earlier (11-13) for investigating the alteration on the level of ChE. Whole brain was homogenized on an ice bath by using a homogenizer in a pH 8.1 barbital-phosphate buffer solution at 3ml/100 mg wet weight of whole blood (11,12). To measure ChE activity, the reaction mixture composed of 3 ml distilled water, 0.2ml plasma or whole brain homogenate and 3ml of pH 8.1 buffer. The initial pH of the mixture (pH₁) was measured by using a pH meter (Hanna, Romania), and then the substrate 7.5 % acetylthiocholine iodide was added to the mixture and incubated at 37°C for 30 min. Then the pH of the reaction mixture (pH₂) should be measured. The ChE activity (ΔpH/30 min.) was calculated as follows:

ChE activity (ΔpH/30 min.)=(pH₁-pH₂)-ΔpH of blank (The blank was without the plasma or brain homogenate sample).

Statistics

To compare the means of two parametric groups, the unpaired Student's t-test was applied for statistical analysis (14,15). On the other hand, the nonparametric data of the two groups were statistically analyzed by using Fisher exact probability test and Mann-Whitney U-test (15,16). The level of significance was at $p < 0.05$.

RESULTS

Acute LD₅₀ value of diazinon in H₂O and H₂O₂ groups of chicks

The acute LD₅₀ value of diazinon was 7.9 mg/kg, orally for both the control and stressed chicks (Table 1). The onset of acute toxicity was decreased and the signs of toxicity (which were salivation, dyspnea, depression, tremor, recumbency, paralysis, convulsion and death) were exacerbated in the stressed group when compared to the control group as demonstrated in Table 1.

Table 1: Acute LD₅₀ value of diazinon in the control and stressed chicks with H₂O₂

Parameters	Results	
	Control group (Tap water)	Stressed group (0.5 % H ₂ O ₂ in water)
Acute LD ₅₀ value	7.9 mg/kg, orally	7.9 mg/kg, orally
The doses range	7-10	7-10
Initial dose	10 mg/kg	10 mg/kg
Last dose	10 mg/kg	10 mg/kg
Increase or decrease in the dose	3 mg	3 mg
Number of chicks	5 (XOXOX)	5 (XOXOX)
Onset of acute toxicity	21-48 min.	5-11 min.
Toxicity signs	Salivation, <u>dyspnea</u> , depression, tremor, <u>recumbency</u> , paralysis, convulsion and death	The same as the control with exacerbation
% effect of OS on acute LD ₅₀ value of <u>diazinon</u> = 0 %		

X= death; O= survival

The ED₅₀ value of atropine in the control and stressed groups of chicks

Table 2 shows that the ED₅₀ value of atropine was effective in treating the diazinon toxicity increased by 265 %, which demonstrates decreased effectiveness in the H₂O₂ (stressed) group (9.5 mg/kg, i.m.) when compared with the H₂O (control) group (2.6 mg/kg, i.m.).

Table 2: ED₅₀ of atropine for preventing the acute toxicity of diazinon in control and stressed chicks with H₂O₂

Parameters	Results	
	Control group (Tap water)	Stressed group (0.5 % H ₂ O ₂ in water)
ED ₅₀ value	2.6 mg/kg, i.m.	9.5 mg/kg, i.m.
The range of the doses	2-4	6-10
Initial dose	4 mg/kg	6 mg/kg
Last dose	3 mg/kg	8 mg/kg
Increase or decrease in the dose	1 mg	2 mg
Number of chicks	6 (XXOOXX)	6 (OOXOXO)
% effect of OS on ED ₅₀ value of Atropine= 265 %		

X= survival; O= death

Atropine was injected (i.m.) immediately after the diazinon administration at 10 mg/5ml/kg, orally

Therapeutic efficacy of atropine against acute diazinon toxicity in control and stressed chicks with H₂O₂

The signs of acute diazinon toxicity were found to increase in the stressed groups when compared to control groups of chicks by measuring the onset of acute signs, signs of toxicity and toxicity scores which revealed also a decrease in atropine therapeutic efficacy in the stressed chicks when compared to the control tap water group of chicks (Table 3).

Table 3: Therapeutic efficacy of atropine against acute diazinon toxicity in control and stressed chicks with H₂O₂

Groups	Onset of toxicity (min.) ⁺	% Toxicity signs						Toxicity scores
		Salivation	Defecation	Tremor	Dyspnea	Recumbency	Paralysis	
Control group (tap water)	49.67 ± 2.88	100	50	16.67	16.67	100	33.33	14
Stressed group (0.5 % H ₂ O ₂ in water)	15.67 ± 2.04*	100	50	66.67*	83.33*	66.67	33.33	18

+ Data represented Mean ± S.E. for 6 chicks/group

* significantly different from the respective control (H₂O group) at p < 0.05

Atropine was injected at 6 mg/5ml/kg, i.m. immediately after diazinon dosing at 10 mg/5ml/kg, orally

ChE activity in the plasma and whole brain of the control and stressed chicks

Table 4 describes a slight change in the plasma and whole brain ChE activity with more decreased in enzyme activity for the stressed chicks when compared to the control chicks.

Table 4: ChE activity in the plasma and whole brain in the control and stressed chicks with H₂O₂

Groups	Plasma		Whole brain	
	ChE activity (Delta pH/30 min.)	% inhibition	ChE activity (Delta pH/30 min.)	% inhibition
Control group (tap water)	0.048 ± 0.005	-----	0.030 ± 0.005	-----
Stressed group (0.5 % H ₂ O ₂ in water)	0.035 ± 0.008	37	0.028 ± 0.009	7

Data represented Mean ± S.E. for 6 chicks/group

Atropine was injected at 6 mg/5ml/kg, i.m. immediately after diazinon dosing at 10 mg/5ml/kg, orally

Blood samples and whole brain were collected after 2 hours of atropine and diazinon administration

DISCUSSION

Many cases of poisoning and diseases in animals may be accompanied by stressful conditions that may affect the pharmacological responses and therapeutic efficacy of certain drugs. The goal of this study was to evaluate the therapeutic efficacy of atropine in case of ChE inhibitors (diazinon as a model here) toxicity in chicks suffered from the OS (as a model of stress) induced with a powerful oxidant H₂O₂ since

no previous literature dealt with that a life threatening situation. Fresh H₂O₂ given each day at 0.5 % in drinking water was found to induce OS in 7-14 day-old chicks old (4,5) which were found by reduced glutathione and elevation of malondialdehyde concentrations in the body tissues of chicks. Atropine is the drug of choice used as an antidote for the treatment of ChE inhibitors (organophosphates and carbamates) toxicity and preventing death in human and animals through blocking the muscarinic receptors (1-3). Diazinon induces toxicity through inhibiting acetylcholinesterase in living tissues (2,3). There was no change in the acute LD₅₀ value for diazinon in both the control and stressed groups of chicks and it may be attributed to acute organophosphate toxicity during 2 hours after dosing and death in vulnerable chicks (17). The therapeutic efficacy of atropine at all was decreased when measuring the ED₅₀ of atropine for treating the diazinon toxicity and death in the control and stressed chicks and this revealed that atropine was not able to treat and counteract diazinon toxicity at the same time with the presence of a stressful condition (OS induced with H₂O₂ as a model) because stressful conditions have synergistic action with toxicity status that exacerbate this condition (18). At the same time, the OS induced with H₂O₂ modulates the pharmacological response and efficacy of drugs in animals (4-7). In the same manner, the signs of acute diazinon toxicity were found to increase in stressed chicks with H₂O₂ when compared to groups that received tap water. A similar effect was described in previous study (18). There was a slight change in the plasma and whole brain ChE activity in stressed chicks when compared to the control chicks which means that the OS status caused by H₂O₂ affect the enzyme level and potentiates the toxicity status (18). The data revealed a decline and modulation in atropine efficacy for treatment of ChE inhibitors (diazinon) toxicity in the

chicks that suffered from OS status induced with H₂O₂. We recommend here an increase the dose of atropine in order to improve the treatment in this case.

الفعالية العلاجية للأتروبين ضد التسمم بمثبطات خميرة الكولين أستراز في الأفراخ المجهدة

بيروكسيد الهيدروجين

يعرب جعفر موسى*؛ منى حازم ابراهيم الزبيدي*؛ سوسن محمد أمين* و فؤاد قاسم محمد**

* فرع الفلسفة والكيمياء الحياتية والأدوية، كلية الطب البيطري، جامعة الموصل، الموصل، العراق.

** وزارة التعليم العالي والبحث العلمي، بغداد، العراق.

الخلاصة

تهدف الدراسة الى تقييم الفعالية العلاجية للأتروبين في حالة التسمم الحاد بمثبطات خميرة الكولين أستراز في نموذج أفراخ الدجاج والتي تعاني من الإجهاد التأكسدي المحدث باستخدام بيروكسيد الهيدروجين حيث أنه لا توجد دراسات سابقة أخذت بنظر الاعتبار هذه الحالة. وقد وجد أن إعطاء بيروكسيد الهيدروجين بتركيز 0.5% في ماء الشرب يعمل على إحداث الاجهاد التأكسدي في أفراخ الدجاج بعمر 7-14 يوم عندما يعطى بصورة طازجة كل يوم. لم يكن هناك تغيير في الجرعة المميتة الوسطية الحادة للديازينون (7,9 ملغم / كغم، عن طريق الفم) لكل من المجموعة التي أعطيت ماء الشرب وتلك التي اعطيت بيروكسيد الهيدروجين مع ماء الشرب. وعند قياس الجرعة الفعالة الوسطية للأتروبين لكل من المجموعة التي أعطيت ماء الشرب (2,6 ملغم / كغم، في العضل) وتلك التي اعطيت بيروكسيد الهيدروجين مع ماء الشرب (9,5 ملغم / كغم، في العضل) فقد تبين أن هناك انخفاض في الفعالية العلاجية (265%) لعلاج التسمم الحاد بالديازينون في أفراخ الدجاج. وعند العلاج بالأتروبين فقد لوحظ زيادة العلامات السمية الحادة للديازينون في المجموعة المعاملة بيروكسيد الهيدروجين بالمقارنة مع المجموعة المعطاة ماء الشرب وفي الوقت نفسه هناك تغييرات طفيفة في نشاط خميرة الكولين أستراز في البلازما والدماغ الكلي الدماغ لكلا المجموعتين. كشفت بيانات البحث عن انخفاض في فعالية الأتروبين العلاجية من التسمم المحدث بمثبطات خميرة الكولين أستراز (الديازينون) في أفراخ الدجاج التي تعاني من الاجهاد التأكسدي المحدث بيروكسيد الهيدروجين وينصح هنا بزيادة جرعة الأتروبين لتعزيز الكفاءة العلاجية من التسمم في هذه الحالة.

REFERENCES

1. Mohammad, F.K., Faris, G.A.-M. and Shindala, M.Kh. (2002). Comparative antidotal effects of diphenhydramine and atropine against dichlorvos–induced acute toxicosis in rats. *Vet. Arhiv.*, 72 (1): 19–28.
2. Taylor, P. (2006). Anticholinesterase Agents. In : Brunton, L.L., Lazo, J.S. and Parker, K.L. (editors), *Goodman and Gilman's the Pharmacological Basis of Therapeutics*. 11th ed., McGraw-Hill Co., Inc., New York, pp. 175–191.
3. Taylor, P., Kovarik, Z., Reiner, E. and Radic, Z. (2007). Acetylcholinesterase: Converting a vulnerable target to a template for antidotes and detection of inhibitor exposure. *Toxicol.*, 233 : 70-78.
4. Mousa, Y.J. (2012). Pharmacological response to some sedatives and analgesics in chicks stressed with hydrogen peroxide. Ph.D. Dissertation, Mosul-Iraq.
5. Mousa, Y.J. and Mohammad, F.K. (2012). Effects of hydrogen peroxide on diazepam and xylazine sedation in chicks. *Interdisc. Toxicol.*, 5(4): 179–183.
6. Mousa, Y.J. (2014). Anaesthetic properties of ketamine in chicks stressed with hydrogen peroxide. *Vet. Med.*, 59 (8): 369–375.
7. Mohammad, F.K., Tawfeek, F.Kh., Hassan, A.A. (1999): Pentobarbital anaesthesia in rats treated with hydrogen peroxide: effect of vitamin E. *Iraqi J. Vet. Sci*, 12: 203–210.
8. Watt, B.E., Proudfoot, A.T., Vale, J.A. (2004): Hydrogen peroxide poisoning. *Toxicol. Rev.*, 23: 51–57.
9. Sayre, L.M., Perry, G., Smith, M.A. (2008): Oxidative stress and neurotoxicity. *Chem. Res. Toxicol.*, 21: 172–188.

10. Dixon, W.J. (1980). Efficient analysis of experimental observations. *Annu. Rev. Pharmacol. Toxicol.*, 20: 441–462.
11. Mohammad, F.K., Faris, G.A.-M. and Al-Kassim, N. A. (1997). A modified electrometric method for measurement of erythrocyte Acetylcholinesterase activity in sheep. *Vet. Hum. Toxicol.*, 39: 337–339.
12. Mohammad, F.K. (2007). Review of a practical electrometric method for determination of blood and tissue cholinesterase activities in animals. *Vet Scan*, 2: 1-12.
13. Al-Badrany, Y.M.A. and Mohammad, F.K. (2007). Effects of acute and repeated oral exposure to the organophosphate insecticide chlorpyrifos on open-field activity in chicks. *Toxicol. Let.*, 174: 110-116.
14. Petrie, A. and Watson, P. (1999). *Statistics for Veterinary and Animal Sciences*. Blackwell Science, Oxford.
15. Katz, M.H. (2006). Bivariate statistics. In: Katz, M.H. (editor), *Study design and statistical analysis*. Cambridge University Press, New York, USA, pp. 66–119.
16. Runyon, R.P. (1977). *Non parametric statistics: A contemporary approach*. Addison–Wesley Publishing Co., Reading, Massachusetts, pp. 2–217.
17. Muna H. I. Al-Zubaidy, Yaareb J. Mousa, Mohammad M. Hasan, and Fouad K. Mohammad (2011). Acute toxicity of veterinary and agricultural formulations of organophosphates dichlorvos and diazinon in chicks. *Arh Hig Rada Toksikol*, 62: 317-323.
18. Al-Baggou, B.K., Naser, A.S. and Mohammad, F.K. (2011). Hydrogen peroxide potentiates organophosphate toxicosis in chicks. *HVM Bioflux*, 3 (2): 142-149.