

## **ANEW METHOD FOR ESTIMATION MEBENDAZOLE IN ITS PHARMACEUTICAL PREPARATIONS AND IN CAMEL URINE**

Nief Rahman Ahmad\* , Farha Khalaf Omar\*\*

\*Department of Environmental Technology, College of Environmental, University of Mosul-Iraq.

\*\*Department of Chemistry, Education College for Girls, University of Mosul-Iraq

(Received 25 June 2018 , Accepted 7 August 2018 )

**Keywords;** Mebendazole, Spectrophotometric method, Camel.


*Corresponding Author: farhakhalf8@gmail.com*

### **ABSTRACT**

A simple, accurate and rapid spectrophotometric method has been developed for the estimation of mebendazole in pure, pharmaceutical formulations and camel urine. The method is based on the reaction of mebendazole with 1-fluoro 2, 4-dinitrobenzene to produce orange-yellow colored complex having maximum absorption at 494 nm. Bee's law was obeyed over the concentration range of 20-280 $\mu$ g/ml, with molar absorptivity of  $3.32 \times 10^4$  l/mol.cm. The present method is considered to be simple because it does not need heating, hydrolysis and solvent extraction steps. The proposed method has been successfully applied for the determination of mebendazole in pure ,pharmaceutical formulations and in camel urine.

### **INTRODUCTION**

Mebendazole, chemically, Methyl-5-benzoyl-2-benzimidazole carbamate (Figure 1)[1], is a benzimidazole derivative that has been widely used in the treatment of hookworm, pinworm, roundworm, tapeworm, threadworm and mixed infestations. It is available in tablet and syrup form. Depending on the type of worm to be treated the dosage varies in adults and children. In Veterinary Medicine it is administered orally to horses at target dose of 8.8mg/kg and to sheep and goats at target dose of 15mg/kg [1-4]. Mebendazole is one of the most widely used drugs for the control of helminthes The drug is known to act through irreversible inhibition of glucose uptake in the parasite, leading to depletion of glycogen store.[5-7].



Methyl-(5-benzoyl-1H-benzimidazol-2-yl) carbamate

C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>3</sub>=295.3

Figure 1: chemical structure of Mebendazole .

Different analytical methods reported for the assay of mebendazole in various dosage forms include infrared spectroscopy [8], differential pulse polarography [9], fluorescence [10], HPLC methods [11,12], and spectrophotometric methods [13,14]. In addition to Liquid chromatographic methods [15], Voltammetric methods [16], volumetric titration method [17], and some UV-spectrophotometric methods [18-20]. The proposed method has been successfully applied for the determination of mebendazole in pure form, pharmaceutical preparations and camel urine.

## MATERIAL AND METHODS

### Apparatus

A spectra scan 50 UV-visible spectrophotometer with 1.0 cm quartz cells was used, [UVS-2700, Labomed, INC], USA.

### Reagents

All chemicals used were of analytical grade and the standard material mebendazole was provided by state company of drug industries and medical appliance (SDI) Samara– Iraq.

### Standard solution of Mebendazole (1 mg/ml):-

0.1 g of mebendazole was dissolved in 10 ml of 1% hydrochloric acid in ethanol, shaken to dissolve and the volume was adjusted to 100 ml with ethanol in a volumetric flask.

### 1-fluoro 2,4-dinitrobenzene(1%):-

This solution was prepared by dissolve 1 g of 1-fluoro 2, 4-dinitrobenzene in 100ml ethanol in a volumetric flask.

**Sodium hydroxide solution (0.1 N):-**

This solution was prepared by dissolving 4 g of sodium hydroxide in 100 ml of distilled water in a volumetric flask.

**Recommended procedure:**

An aliquots of standard solution of Mebendazole (0.5-7.0 mg) were transferred into a series of 25ml volumetric flasks, 1 ml of 1N NaOH and 2 ml of 1-fluoro 2, 4-dinitrobenzene solution were added. The contents were diluted to the mark with distilled water. The absorbance was measured at 488 nm against a reagent blank.

**Procedures for pharmaceutical preparations (tablets):**

To minimize a possible variation in the composition of the tablets (100mg of Mebendazole /tablet)was used. Ten tablets were weighed , grounded, and powdered . Mebendazole (100mg) was then accurately weighed and added 10 ml of 1% hydrochloric acid in ethanol, shake for 30 minutes and diluted to 100.0 ml with ethanol and filtered using a whatman No.42 filter paper . Different volume of this solution was treated as mentioned under recommended procedure.

**Procedures for pharmaceutical preparations (suspensions):**


The contents of five bottles of suspensions.5ml of Vermosam suspension contain 100mg of Mebendazole) were mixed well. An aliquot corresponding to 100mg (5ml) of suspensions was diluted to 100 ml with ethanol in a volumetric flask. Different volume of this solution was treated as mentioned under recommended procedure.

**Procedure for spiked camel urine: -**

To demonstrate the practical applicability of the proposed method, 10 ml of camel urine were analyzed by this method. Camel urine was fortified with 1, 4 and 6 ml of 0.1% Mebendazole solution. The fortified camel urine samples were analyzed as described above under recommended procedure.


## **RESULTS AND DISCUSSION**

The reaction of nucleophile substitution between mebendazole and 1-fluoro 2, 4-dinitrobenzene with the purpose of elaborating a new spectrophotometric method for the determination of mebendazole was investigated. The reaction occurred in slightly alkaline medium produces an orange-yellow color with maximum absorbance at 494 nm (Figure 2).



**Figure 2: Absorption spectra of 3mg/25ml of mebendazole with 1-fluoro 2, 4-dinitrobenzene against reagent blank(A). Reagent blank against water (B).**

The reaction variables were optimized by varying each variable and keeping others constant for obtaining maximum absorbance. The reaction was found to be quantitative in the sodium hydroxide medium. It was found that 1 ml of 1 N sodium hydroxide solution gives high sensitivity, 1-fluoro 2, 4-dinitrobenzene on the absorbance was investigated. A maximum and constant absorbance was founded with 1 to 2 ml of 1% of 1-fluoro 2, 4-dinitrobenzene solution and 2.0 ml has been used for subsequent experiments. The color reaction occurred at room temperature immediately and remained stable for at least 24h and a reaction time of 5 min was selected for reproducible results. Under the experimental conditions described, Beer's law is obeyed over the concentration range 20-280 $\mu$ g/ml (Figure 3).



**Figure 3: Calibration curve of Mebendazole.**

A regression analysis of Beer's law plot at 488 nm revealed a good correlation ( $r=0.9999$ ,  $n=7$ ). The graph of the absorbance versus the concentration of mebendazole showed a low intercept ( $-0.017$ ) and slope ( $0.028$ ) described by a regression equation  $Y = ax + b$  (where  $x$  is the concentration of mebendazole in  $\mu\text{g}/\text{ml}$ ,  $Y$  is the absorbance,  $a$  is the slope and  $b$  is the intercept). The apparent molar absorptivity was  $1.346 \times 10^4 \text{ l. mol}^{-1} \cdot \text{cm}^{-1}$ . The optical characteristics are given in (Table 1).

**Table 1: Optical characteristics and statistical data for regression equation**

Parameters	Value
$\lambda$ max (nm)	488
Beer's law limits ( $\mu\text{g} \cdot \text{ml}^{-1}$ )	20 – 280
Molar absorptivity ( $1 \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$ )	$3.32 \times 10^4$
Determination coefficient ( $r^2$ )	0.9987
Regression equation ( $Y = a \times + b$ )	
Slope (a)	0.0785
Intercept (b)	0.0077

**Accuracy and precision**

The accuracy and precision of the method was established by analyzing the pure drug solution at three different levels. The average recovery which is a measure of accuracy is  $100 \pm 1.5$  revealing high accuracy of the method. The relative standard deviation (RSD), which is an indicator of precision is better than  $\pm 2\%$  (Table 2).


**Table 2: Accuracy and precision of the proposed method.**

Mebendazole taken $\mu\text{g}/\text{ml}$	Mebendazole found $\mu\text{g}/\text{ml}$	Er (%) <sup>a</sup>	RSD (%)
20	20.3	1.5	1.9
100	99.4	0.6	1.3
200	201.5	0.75	1.5

a: Mean of six determinations

**Stoichiometry of reaction**

The stoichiometry of the reaction between mebendazole and 1-fluoro 2, 4-dinitrobenzene was investigated using job's method of continuous variation and mole ratio methods of equal molar solution ( $3.386 \times 10^{-3} \text{M}$ ). The result obtained showed that (1:1) 1-fluoro 2, 4-dinitrobenzene to drug at 488 nm. The suggested reaction and structure of the product might be written as:



### Apparent stability of the product

The conditional stability constant of the product can be estimated by using the following equation [21]

$$K = a - (\Delta A / \epsilon) / n^n (\Delta A / \epsilon)^{n+1} \quad \text{Where: } a = \text{total mebendazole concentration. (Molar)}$$

$\Delta A$  = Sample absorbance in reagent excess minus the sample absorbance at stoichiometric amount.

$\epsilon$  = Molar absorptivity at the measured wavelength and  $n$  = number of ligand .

The stability constant (means of three values) was found to be  $1.48 \times 10^7$  l/mol indicating the product is very stable.

### The effect of foreign compounds or excipients in assay of mebendazole

The interfering effect of foreign species often accompanied with mebendazole in the pharmaceutical preparations were studied by adding different amounts of foreign species to  $200 \mu\text{g/ml}$  of mebendazole in solution .The recommended procedure for the determination of mebendazole was followed . The species are considered to interfere seriously if the cause aching of more than 2% in the absorbance obtained for mebendazole alone [22]. The results of the

recovery analysis are presented in (Table 3). Excipients at the concentration revealed do not interfere with the assay (Table 3).In addition recoveries in most cases were around 100%.

**Table 3: Determination of mebendazole in presence of excipients**

<b>Excipients</b>	<b>Amount taken, (µg/ml)</b>	<b>Average recovery, * %</b>
<b>Talc</b>	<b>1000</b>	<b>100.10</b>
<b>Mannitol</b>	<b>1000</b>	<b>100.16</b>
<b>Mg – stearate</b>	<b>1000</b>	<b>100.12</b>
<b>Starch</b>	<b>1000</b>	<b>99.90</b>
<b>Microcrystalline cellulose</b>	<b>1000</b>	<b>99.98</b>

\* Average of three replicate determinations.

**Application of the proposed method**

The proposed method was successfully applied to the analysis of mebendazole in Vermosam tablets, Vermosam suspension, and spiked camel urine samples. The result of analysis for pharmaceutical formulations revealed that there is close agreement between the results obtained



by the proposed method and the label claim (Table 4), and the results of spiked camel urine samples. The recovery values obtained were close to 100% (Table 5).

**Table 4: Assay of mebendazole in pharmaceutical formulations.**

Pharmaceutical formulation supplied by SDI	Amount of mebendazole*	Label claim	%Recovery
	Proposed method		
Vermosam tablets 100mg/tablet	99.4 mg	100 mg	99.4
Vermosam suspension [each 5ml contain 100mg]	100.1mg/5ml	100mg/5ml	100.1

\*Mean of ten determinations.

**Table 5: Determination of mebendazole in spiked camel urine**

samples	mebendazole( $\mu\text{g/ml}$ )*		%Recovery
	Taken	Found	
camel urine	40	40	100
	100	99.5	99.5
	160	159	99.37

\*Mean of ten determinations.

## CONCLUSION

In the current study, a simple, rapid, precise and accurate spectrophotometric method was developed and validated for the determination of mebendazole in pharmaceutical preparations and spiked camel urine samples. The method free from such experimental variables as heating or solvent extraction step. The method rely on the use of simple and cheap chemicals and techniques and can be used for rapid routine determination and quality control of mebendazole in pure form, pharmaceutical preparations (tablets, suspension) and spiked camel urine sample.

## ACKNOWLEDGMENTS

The authors wishes to express gratitude to [the state company of drug industries and medical appliance (SDI) Samara– Iraq for providing gift samples of mebendazole standard materials and pharmaceutical preparations (tablets and suspension).

### طريقة جديدة مبتكرة لتقدير الميبندازول في مستحضراته الصيدلانية وفي ادرار الجمل

نايف رحمن احمد، فرحة خلف عمر

### الخلاصة

تم اختبار طريقة طيفية جديدة تمتاز بالبساطة والدقة والسرعة لتقدير ميبندازول في حالته النقية وفي بعض مستحضراته الدوائية وفي ادرار الجمل، تعتمد الطريقة على تفاعل ميبندازول مع ١- فلور ٢,٤- ثنائي نايترو بنزين لتعطي ناتج برتقالي- اصفر اللون له أقصى امتصاص عند الطول الموجي ٤٩٤ نانومتر وأمكن تقدير الكميات التي تتراوح بين ٢٠-٢٨٠ مايكروغرام/مل وان معامل الامتصاص المولاري للطريقة المقترحة هو  $3.32 \times 10^4$  لتر.مول<sup>-١</sup>.سم<sup>-١</sup>. تتميز الطريقة بالبساطة والانتقائية والسرعة والدقة وتعتبر بسيطة كونها لا تحتاج للتسخين ولا للاستخلاص المذيب. وتم دراسة الظروف المثلى للتفاعل. وطبقت الطريقة بنجاح لتقدير ميبندازول في حالته النقية وفي مستحضراته الصيدلانية وفي ادرار الجمل.

## REFERENCES

- 1- British pharmacopeia, Her Majesty, Stationary office, London,(2014), Volume2, p.193.
- 2- United States Pharmacopeia and National Formulary,(2018), USP 41, NF 36 .P 2534-2536.
- 3- Martindale "The Extra Pharmacopeia -The Complete Drug Reference" edited by Sean. C, Sweetman, Pharmaceutical press, London, UK,(2009), 36th ed. P.148-149
- 4- British National Formulary (BNF) Royal Pharmaceutical Society.(2016), 70, p.526
- 5-Leveck B, Montresor A, Albonico M, Ame SM, Behnke JM (2014), 'Assessment of Anthelmintic Efficacy of Mebendazole in School Children in Six Countries Where Soil-Transmitted Helminths Are Endemic. PLo SNegl Trop Dis, 8, 3204.

- 6- **Zeugin, T., Zysset, T. and Cotting, J. (1990),**'Therapeutic monitoring of albendazole: A high-performance liquid chromatography method for determination of its active metabolite albendazole sulfoxide. *Therapeutic Drug Monitoring*, 12,187-190
- 7- **Pinto, L.C., Soares, B.M., Pinheiro, J.J., Riggins, G.J., Assumpção, P.P., Burbano, R.M., Montenegro, R.C.(2015),**'The anthelmintic drug mebendazole inhibits growth, migration and invasion in gastric cancer cell model., *Toxicol. in Vitro*, 29, 2038–2044
- 8-**Gunasekaran .S and Uththra. D,(2008),**'Vibrational Spectra and Qualitative Analysis of Albendazole and Mebendazole, *Asian Journal of Chemistry*, 20(8), 6310-6324
- 9- **Christine Jeyaseelan and A P Joshi,(2009),**Study of Mebendazole by Differential Pulse Polarography, *International Journal of ChemTech Research*, 1(3), 596-601.
- 10- **Baeyens.W, Abdel Fattah.F, DeMoerloose .P(1985),** 'Fluorimetric analysis of mebendazole and flubendazole with hydrogen peroxide; *J. Pharm. Biomed. Anal.* 3, 397-404
- 11-**Anna K, Wanda B, Marzena P, Barbara K and Aleksander M,(2011) ,** ' HPLC Method for identification and quantification of benzimidazole derivatives in antiparasitic drugs, *Acta Poloniae Pharmaceutica n Drug Research*,68(6), 823-829.
- 12-**Turabi Z.M.,Khatatbeh O.A,Al-Abed D.N ,(2014);** RP-HPLC Method Development and Validation for the Simultaneous Determination of mebendazole and the two preservatives methylparaben and propylparaben in Pharmaceutical Oral suspension dosage form; *pharm. sciences and drug research*, 6, 70-74.
- 13- **Swamy, N, Prashanth, K, Basavaiah, K,(2014),**'Spectrophotometric Assay of Mebendazole in Dosage Forms Using Sodium Hypochlorite. *Journal of Applied Spectroscopy* . 81 (3), 525-533.
- 14- **Swamy. N, Basavaiah.K.(2013),**' Selective and sensitive assay of mebendazole in pharmaceuticals using bromocresol green by spectrophotometry, *Thai J. Pharm. Sci* ,37, 171-185.
- 15- **Ruyck Hendrik DE, Daeseleire Els, Ridder Herman DE, and Vanrenterghem Roland,(2003),**' Liquid chromatographic-electro spray tandem mass spectrometric method for the determination of Mebendazole and its hydrolysed and reduced metabolites in sheep muscle, *Analytica chemical Acta*, 483(1-2), 111-123.

- 16- Lopes D, Fereira G, Ribeiro S, Lima S, Fernandez C, Goncalves S, Souza M, Sapateiro Tand De Souza.(2016),**'Differential pulse voltammetric determination of albendazole and mebendazole in pharmaceutical formulations based on modified sonogel carbon paste electrodes with Perovskite-type LaFeO<sub>3</sub> nanoparticles. Journal of the Electrochemical Society [online], 163(8) , B428-B434.
- 17- Raja C, Sampath K, Jayakumar C, Nagendra G. N.(2013),**' Quantitative analysis of bendazole bulk sample using sodium salicylate hydrotropic agent, Int J Curr Pharm Res, 5(2), 93-94
- 18- Madkour.(2015),**'Spectrophotometric Methods for Determination of Mebendazole in Presence of its Alkaline Induced Degradation Product in Pure Form and Pharmaceutical Preparation, Ijppr.Human, 4 (3), 1-19.
- 19--Durgesh Parakh,(2015),**' Development and validation of spectrometric method for estimation of mebendazole in bulk and pharmaceutical formulation, World Journal of Pharmaceutical Research, 4(7),2223-2235
- 20- Mario R. D, Celina M. M, Nelly L. J and María del C. S,(2017),**'Mebendazole spectrophotometric determination ,theoretical and experimental study of the interaction with sodium hydroxide, Asian Journal of Science and Technology,7 (11), 3948-3953.
- 21- Nief.Rahman Ahmad and Widad Aesa.(2009),**'Spectrophotometric determination of Chlorocresol in some pharmaceutical preparations through nitrosation reaction, J.Raf.Sci, 20(3), 66-73
- 22- Nief Rahman Ahmad.(2012),**' Facial visible spectrophotometric determination of metformin Hydrochloride in glucosam tablets and industrial waste water:Application to content uniformity testing, Iraq J Pharm,12(1),75-85.